



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



$VD$  cutting  $VC$  at  $C$ . Produce  $CD$  to  $P$  making  $DC = DP$ . Draw the lines  $AP$ ,  $BP$ , and  $VP$ . From this construction it easily follows that  $\angle BVC = \angle PVA$  and  $\angle CVP = 2\angle CVD$ . But  $\angle CVP < \angle AVC + \angle PVA$ . Hence  $\angle CVD < \frac{1}{2}(\angle AVC + \angle BVC)$ .

(2)  $\angle CVD > 90^\circ$ . (Fig. 2.) Produce  $CV$  through  $V$  and draw plane  $ABC'$  as in (1). Then  $2\angle C'VD < \angle AVC' + \angle BVC'$  by (1). Hence  $2(180^\circ - \angle C'VD) > 180^\circ - \angle AVC' + 180^\circ - \angle BVC'$  or  $\angle CVD > \frac{1}{2}(\angle AVC + \angle BVC)$ .

(3)  $\angle CVD = 90^\circ$ . (Fig. 3.) In this case the plane through  $AB$  is parallel to  $VC$ . Draw a plane  $MNV$  through  $VD$  perpendicular to  $VC$  and cut by planes  $BVC$  and  $AVC$  in the lines  $MV$  and  $NV$  respectively. It can be easily shown that  $\angle MVB = \angle NVA$ . Hence

$$\angle AVC + \angle BVC = 180^\circ$$

and

$$\angle CVD = \frac{1}{2}(\angle AVC + \angle BVC).$$

Also solved by B. LIBBY, F. M. MORGAN, and GEO. W. HARTWELL.

#### CALCULUS.

##### 338. Proposed by RICHARD LOCHNER, Philadelphia, Pa.

An elliptical field has a major axis of 100 feet and a minor axis of 10 feet. A cow is tethered at the end of the major axis and another at the end of the minor axis. If each cow can graze over half the field, how long is the rope of each? What is the area of the portion over which the cows can graze in common?

SOLUTION BY B. F. FINKEL, Drury College.

The central equation of the elliptic field is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

Let  $r_1$  be the length of rope by which one cow is tethered at the point  $A_1$ , the right-hand extremity of the major axis. The equation of the circle over which this cow can browse is  $(x - a)^2 + y^2 = r^2$ . The coördinates of the point of intersection,  $P_1$ , of this circle with the ellipse are

$$(x_1, y_1) \equiv \left( a \frac{[a^2 - \sqrt{b^4 + r^2(a^2 - b^2)}]}{(a^2 - b^2)}, \right. \\ \left. \sqrt{2a^4 \frac{\sqrt{b^4 + r^2(a^2 - b^2)} - b^2(a^2 - b^2)r^2 - a^2(a^4 + b^4)}{(a^2 - b^2)}} \right),$$

The area common to the ellipse and the circle is

$$\text{area} = 2 \int_0^{y_1} \int_{a - \sqrt{r^2 - y^2}}^{(a/b)\sqrt{b^2 - y^2}} dx dy \\ = 2 \left[ \frac{a}{b} \frac{y_1}{2} \sqrt{b^2 - y_1^2} + \frac{b^2}{2} \sin^{-1} \frac{y'}{b} - ay' + \frac{y'}{2} \sqrt{r^2 - y_1^2} + \frac{r^2}{2} \sin^{-1} \frac{y_1}{r} \right],$$

which, by the conditions of the problem,  $= \frac{1}{2}\pi ab$ .